arXiv:1704.03485v1 [math.KT] 11 Apr 2017

This paper is dedicated to the 80-th anniversary of Yuri I. Manin

SUBLINEAR EXTENSION OF GROTHENDIECK ALGEBRAIC K–THEORY

IGOR V. ORLOV

Key words: Grothendieck group, algebraic K–theory, divisible Abelian semigroup, uniquely divisible Abelian semigroup, convex cone, linear space, cancellation law, formal difference, canonical embedding.

AMS Mathematics Subject Classification: Primary 16E20, 18F25, Secondary 49J52.

Abstract. A commutative diagram that connects the basic objects of commutative algebra with the main objects of commutative analysis is constructed. Namely, with the help of five types of canonical embeddings we constructed a diagram between two sets of objects: Abelian semigroups – Abelian regular (cancellative) semigroups – Abelian groups, on the first hand, and convex cones – regular convex cones – linear spaces, on the other hand. Thus, some extension of the Grothendieck algebraic K–theory arises, that includes the basic objects not only of linear (smooth) analysis but of sublinear (nonsmooth) analysis also.

1. Basic objects: terminology, notation, auxiliary results

(1) *Abelian semigroup* with zero element (i.e., monoid): $X = \{x\}$, with additive notation. The corresponding category denote by (S).
(2) *Regular (cancellative) Abelian semigroup* is an Abelian semigroup that satisfies the *cancellation law*: $(x + y = x + z) \Rightarrow (y = z)$ (see [1]). The corresponding category denote by (RS).
(3) *Abelian group*. The corresponding category denote by (G).
(4) *Convex cone* is an Abelian semigroup with respect to vector addition that forms a module over $\mathbb{R}_+$ with respect to multiplication by scalars (see [2, 3]). The corresponding category denote by (Con).

(5) *Regular (cancellative) convex cone* is a convex cone that satisfies the cancellation law (see [2, 3]). The corresponding category denote by $(RCon)$.
(6) *Linear space* (over $\mathbb{R}$). The corresponding category denote by (Lin).

Let's remind some auxiliary concepts and results.

Proposition 1.1. ([1]) *An arbitrary Abelian semigroup can be isomorphically (injectively and additively) embedded into some Abelian group if and only if it is regular.*

Corollary 1.2. ([2]) *An arbitrary convex cone can be isomorphically (i.e. injectively and $\mathbb{R}_+-$linearly) embedded into some linear space if and only if it is regular.*

Definition 1.3. ([4, 5, 6]) The minimal Abelian group that contains given regular semigroup X is called *Grothendieck group* of X and is denoted by $Gr(X)$. Respectively, the minimal linear space that contains given regular convex cone X let's call *Grothendieck linear group* of X and is denoted by $Gr_L(X)$.

2. Divisible objects: terminology, notation, auxiliary results

(1) *Divisible Abelian semigroup* is an Abelian semigroup X that satisfies the condition:

$$\forall\, x \in X\ \forall\, n \in \mathbb{N}\ \exists\, y \in X : \underbrace{y + \cdots + y}_{n} := \sum\nolimits_n y = x$$

(see [7]). The corresponding categories denote by (DS), (DRS), (DG).

(2) Uniquely divisible Abelian semigroup is a divisible Abelian semigroup X that satisfies the condition:

$$\forall\, x_1, x_2 \in X,\ \forall\, n \in \mathbb{N}\ \left(\sum\nolimits_n x_1 = \sum\nolimits_n x_2\right) \Rightarrow (x_1 = x_2)$$

(see [7, 8]). The corresponding categories denote by (US), (URS), (UG).

Proposition 2.1. *Let X be a uniquely divisible Abelian semigroup. Then:*

$$\forall\, x \in X,\ \forall\, n_1, n_2 \in \mathbb{N} \quad \left(\sum\nolimits_{n_1} x = \sum\nolimits_{n_2} x,\ x \neq 0\right) \Rightarrow (n_1 = n_2).$$

3. Basic canonical embeddings

Let's describe in short the main embeddings that will be used further.

(1) *Regularization.* $R : (S) \to (RS)$, $(DS) \to (DRS)$, $(US) \to (URS)$, $(Con) \to (RCon)$.

Factorize X by $(y_1 R y_2) \Leftrightarrow (\exists\, x \in X : x + y_1 = x + y_2)$.

Let $X_R = X/R$ be corresponding factor semigroup, then $R : X \to X_R$, $y \mapsto \{z \in X |\ yRz\}$ is the required embedding and $X_R \in (RS)$.

(2) *Formal difference.* $F : (RS) \to (G)$, $(DRS) \to (DG)$, $(URS) \to (UG)$, $(RCon) \to (Lin)$.

Factorize $X \times X$ by

$$((y_1, z_1) F (y_2, z_2)) \Leftrightarrow (y_1 + z_2 = y_2 + z_1)\,.$$

Let $X_F = X \times X/F$ be corresponding factor semigroup. Introduce the subtraction operation in X_F by involution

$$-(y, z) = (z, y).$$

Then $F : X \to X_F$ $\left(x \mapsto \{(y, z)\big|\ x + y = z\}\right)$ is the required embedding and $X_F = F(X) - F(X)$.

(3) *Divisibility.* $D : (S) \to (DS)$, $(RS) \to (DRG)$, $(G) \to (DG)$.

Factorize $X \times \mathbb{N}$ by

$$((x_1, n_1) D (x_2, n_2)) \Leftrightarrow \left(\sum\nolimits_{n_2} x_1 = \sum\nolimits_{n_1} x_2\right).$$

Let $X_D = X \times \mathbb{N}/D$ be corresponding factor semigroup, then $D : X \to X_D$ $\left(x \mapsto \{(y, n)\big|\ (x, 1) D (y, n)\}\right)$ is the required embedding.

(4) *Uniquely divisibility.* $U : (DS) \to (US)$, $(DRS) \to (URG)$, $(DG) \to (UG)$.

Factorize $X =$ by $(x_1 U x_2) \Leftrightarrow (\exists\, n \in \mathbb{N} :\ \sum_n x_1 = \sum_n x_2)\,.$

Let $X_U = X/U$ be corresponding factor semigroup, then $U : X \to X_U$ $(x \mapsto \{y \in X |\, \exists\, n \in \mathbb{N} :\ \sum_n x = \sum_n y\})$ is the required embedding.

(5) *Modulation.* (see [3]) $M : (US) \to (Con)$, $(URS) \to (RCon)$, $(UG) \to (Lin)$.

Let's introduce in uniquely divisible Abelian semigroup X an "additive multiplication" by non-negative scalars, first for rational case.

(a) For $x \in X$, $r = \frac{m}{n} \in \mathbb{Q}_+$ set $(y = r * x) \Leftrightarrow (\sum_m x = \sum_n y)$.

(b) For $\gamma \in \mathbb{R}_+$ that defined by Dedekind cutting $A|B$ in $\mathbb{Q}_+$, set

$$\gamma * x = (A * x \big| B * x).$$

(c) Define X_M as additive envelope of the set $\mathbb{R}_+ * X$ with respect to Minkowsky addition:

$$X_M = \left\{ \sum_{k=1}^{n} \gamma_k * x_k \big| \, \gamma_k \in \mathbb{R}_+, \ x_k \in X, \ n \in \mathbb{N} \right\}.$$

Here the modulation is extended to X by obvious way:

$$\alpha * (\sum_{k=1}^{n} \gamma_k * x_k) = \sum_{k=1}^{n} (\alpha\gamma_k) * x_k.$$

The canonical embedding $M : X \to X_M$ is defined by the equality:

$$Mx = 1 * x = \left([0,1]_{\mathbb{Q}} * x \big| (1, +\infty)_{\mathbb{Q}} * x \right).$$

4. Properties of the basic embeddings

The following statements can be checked by direct transformations.

Theorem 4.1. *The embedding R is an additive homomorphism from X onto X_R.*

Theorem 4.2. *The embedding F is an additive isomorphism from X into X_F. In addition, in case $(RCon) \to (Lin)$, F is $\mathbb{R}_+$-linear isomorphism.*

Theorem 4.3. *The embedding D is an additive isomorphism from X into X_D. In addition, $X_D = D(X) - D(X)$.*

Theorem 4.4. *The embedding U is an additive homomorphism from X onto X_U.*

Theorem 4.5. *The embedding M is an additive isomorphism from X into X_M. In addition, $X_M = Add(\mathbb{R}_+ * X)$.*

5. The main result

Theorem 5.1. *The following diagram is commutative, together with any its subdiagram.*



Final Remark. The red arrows on the diagram below show a connection between Theorem 5.1 and classical Grothendieck Theorem (see [4],[5],[6],[9],[10]).

References

[1] A. H. Clifford, G. B. Preston *The Algebraic Theory of Semigroups*, Math. Surveys **7** (1961) vol.1, Math. Surveys **7** (1962) vol.2.

[2] E. S. Polovinkin, M. V. Balashov *Elements of Convex and Strongly Convex Analysis*. Moscow, PhysMathLit Publ (2004) (In Russian).

[3] I. V. Orlov *On Embedding of Uniquely Divisible Abelian Semigroup Into Convex Cone* Math. Notes (2017) (to appear).

[4] Yuri I. Manin *Lectures on the K–functor in algebraic geometry*, Russian Math. Surveys **24**:5 (1969) 1–89.

[5] M. F. Atiyah *K–theory*. W. A. Benjamin Inc., New York (1967).

[6] P. N. Achar, C. Stroppel *Completions of Grothendieck Groups*, Bull. of the London Math. Soc. **45** (1) (2013) 200–212.

[7] P. A. Griffith *Infinite Abelian Group Theory*, Chicago Lecture in Math., Univ. of Chicago Press (1970).

[8] J. Tabor *k–Proper Families and Almost Approximately Polynomial Functions*, Glasnik Mathematički **36** (56) (2001) 177–191.

[9] M. Karoubi *K–Theory. An Elementary Introduction* arXiv:math/0602082 (2006).

[10] Ch. Weibel *The K–Book. An Introduction to Algebraic K–Theory* Grad. Stud. in Math. **145** AMS (2013).

Igor Vladimirovich Orlov
Department of Mathematics and Informatics
Crimean Federal V. Vernadsky University
Academician Vernadsky Ave., 4
Simferopol, Republic Crimea, Russia, 295007
E-mail: igor_v_orlov@mail.ru